Discovery of the “Compton Shoulder” in the Iron Line from GX301–2

Shin Watanabe(ISAS)

M. Sako, M. Ishida, Y. Ishisaki,
S. Kahn, T. Kohmura. U. Morita,
F. Nagase, F. Paerels, T. Takahashi

Introduction(Compton scattering)

When an X-ray photon (E_0) is Compton-scattered by an electron in a cold material,

$$E_1 = \frac{E_0}{1 + E_0 / mc^2 (1 - \cos\theta)}$$



If any X-ray emission lines are observed,

"Compton shoulder"

- the direct evidence of Compton scattering
- a probe for the physical states of the materials

Observations with high energy resolution hold the key!

Iron Kα line(6.4keV) -> Compton shoulder: 6.24-6.40keV
width : 160eV

Chandra HETG is capable of detecting Compton shoulders

Observation of GX301–2 with Chandra HETG



GX301–2

- High Mass X-ray Binary
  (NS and super-giant(B2Iae))
- Strong iron Kα line (EW ~ 1keV)
  –> the most promising target!!

We have observed at 3 orbital phases
(Intermediate(0.167-0.179) ~40ksec
Near-Apastron(0.480-0.497) ~60ksec
Pre-Periastron(0.970-0.982) ~40ksec)

Result of the observation

Energy spectrum from HEG ±1 order data



A “clear” Compton shoulder is detected for the first time!

Variability of the Compton shoulder

Light curve of HEG ±1 order



X-ray outburst is seen in the first half.

We divided the data into the first and second halves



The profile of the Compton shoulder is changed !

The first ~20 ksec The second ~20 ksec

How to reproduce the shoulder

Issues to be considered

- Two competing physical processes
  (photoelectric absorption and Compton scattering)
- Multiple scatterings
  $N_H \sim 10^{24} cm^{-2}$ –> Optical depth of each process is ~1
- Geometry



It is too complex for analytical approaches.

The easiest way is the Monte Carlo simulation of photon transport.

The Monte Carlo simulation

- A homogeneous spherical cloud
  - The NS (the power-law X-ray source) is located at the center
- Photoelectric absorption and Compton scattering are implemented



Results of Monte Carlo simulations(1)

A transition when N_H is changed. (kT_e is fixed 0eV)

(Metal Abundance: 0.7 cosmic)





The flux ratio of the shoulder to the line vs N_H

Metal abundance
for cosmic abundance(Feldman1992)

0.5 x cosmic

0.7 x cosmic

1.0 x cosmic

Results of Monte Carlo simulations(2)

A transition when kT_e is changed. (N_H is fixed 1×10^{24} cm^{-2})

(Metal Abundance: 0.7 cosmic)



The flux ratio of the shoulder to the line

→ the optical depth and the abundance of the cloud

The shape of the shoulder

→ the temperature of scattering electrons

Fitting of the Compton shoulder



0.7 cosmic abundance is adopted as the metal abundance

N_H 11.7+2.3/-2.0 x10^{23} cm^{-2}, 8.4±1.6 x10^{23} cm^{-2}

Upper limit of kTe < 3.7eV, <0.7eV (90%)

The new means for probing materials around X-ray sources!

N_H from the Compton shoulder can reproduce the whole spectrum shape

The first half



The second half



Full Monte Carlo simulation

- a spherical cloud
- a power-law source with Γ=1.0
- a metal abundance: 0.7 cosmic
- N_H derived from the Compton shoulder

We can obtain a self-consistent solution from the analysis of the Compton shoulder !

Summary

- We discovered the Compton shoulder in the iron Kα line of GX301–2
- From properties of the Compton shoulder, we have obtained physical parameters characterizing the material around the X-ray source, using Monte Carlo simulations.
- The new means for probing cold materials using Compton shoulders was established.

Future

ASTRO-E2 XRS

- Larger effective area(300cm^2@6keV)
  –> other X-ray binaries, AGNs
  diffuse sources (the Galactic center region)
- Excellent energy resolution (6eV@6keV)
  –> geometrical conditions